



柏際股份有限公司 **BockyTech, Inc.**

100台北市中正區延平南路70號2F之3

Tel : 886-2-23618050 Fax : 886-2-23619803

2F-3, No 70, Yanping S. Rd.,
Taipei 100, Taiwan, R.O.C.
[Http://www.bockytech.com.tw](http://www.bockytech.com.tw)

Conners' Continuous Performance Test II (CPT II) Ver. 5.2 for Windows®

- C. Keith Conners, Ph.D. and MHS Staff with additional comments by IPS Educational Psychologist,
Jenny Lyon

Important additional features of CPT II Ver. 5.2 compared with earlier versions:

- The introduction of a short practice exercise for clients prior to the administrations of the full test.
- The option to minimise false positives or false negatives.
- A Confidence Index adjustment to confirm the validity of the administration.
- The Multi-Modal Integrated report combines results from the CPT with the Conners' Rating Scales.
- Activation Key by USB flash drive.
- Compatible with Windows Vista.

Aim

Conners' Continuous Performance Test (CPT II) Version 5.2 for Windows® is an attention test that is widely used in ADHD research and clinical assessments for respondents aged 6 or older. Response patterns on the CPT II provide information that enables the practitioner to better understand the type of deficits that might be present. For example, some response patterns suggest inattentiveness or impulsivity, while other response patterns may indicate activation/arousal problems or difficulties maintaining vigilance.

Potential users of the CPT II include psychologists, social workers, physicians, counsellors, psychiatric workers, paediatric nurses, teachers and school officials. Appropriate settings for the CPT II include schools, outpatient clinics, inpatient clinics, residential treatment centres, child protective services for both placement and referral decisions, special education and regular classrooms, juvenile detention centres, and private practice offices (psychological, psychiatric, paediatric and family practice). The program is commonly used as a screening tool to identify potential attention problems, and as an aid in monitoring treatment effectiveness.